







FOUR REPORTS OF PREDATION OF *Quiscalus mexicanus* (PASSERIFORMES: ICTERIDAE) IN PANAMA CITY.

**¹Ramy Jhasser Martínez J, ^{2,3,4}Jorge Moisés Herrera, Venicio Wilson⁵,
⁶Jacobo Araúz G, ^{1,6}Dora Isabel Quirós**

¹Laboratorio de Ecología y Biodiversidad Neotropical, Vicerrectoría de Investigación y Postgrado, Universidad de Panamá. Email: ramymartonez1009@gmail.com 

²Instituto Smithsonian de Investigaciones Tropicales (STRI), apartado 0843-03092, Balboa, Ancón, República de Panamá. ³Comisión Panamá – Estados Unidos para la Erradicación y Prevención del Gusano Barrenador del Ganado (COPEG). Pacora Centro, Antiguo Ingenio Felipillo, Pacora, República de Panamá. ⁴Sociedad Mesoamericana de Biología y Conservación. ⁵Sociedad Audubon de Panamá. 

⁶Departamento de Zoología, Facultad de Ciencias Naturales Exactas y Tecnología, Universidad de Panamá. 

¹Laboratorio de Ecología y Biodiversidad Neotropical, Vicerrectoría de Investigación y Postgrado, Universidad de Panamá. ⁶Departamento de Zoología, Facultad de Ciencias Naturales Exactas y Tecnología, Universidad de Panamá. 

ABSTRACT

We report the predation of two species of rodents of the Muridae family (*Rattus norvegicus* and *Mus musculus*), one specimen of *Columbina talpacoti* (Aves: Columbiforme) and a crustacean *Leptuca* sp by females of *Quiscalus mexicanus* in Panama. One observation was made in 2008, and four in 2020 in Panama City. This report seeks to promote research on the changes that *Q. mexicanus* makes in its diet depending on the availability of nutritional resources and thus expand the knowledge about its food sources.

KEY WORDS: anthropogenic landscape, bird ecology, dove, eating behavior, rodents, urban area.

RESUMEN

Reportamos la depredación de dos especies de roedores de la familia Muridae (*Rattus norvegicus* y *Mus musculus*), un espécimen de *Columbina talpacoti* (Aves: Columbiformes) y un crustáceo *Leptuca* sp por hembras de *Quiscalus mexicanus* en Panamá. Una observación fue hecha en 2008, y cuatro en 2020 en la Ciudad de Panamá. Este reporte busca promover la investigación sobre los cambios que hace *Q. mexicanus* en su dieta dependiendo de la disponibilidad de recursos alimenticios y así expandir el conocimiento acerca de sus fuentes de alimento.

PALABRAS CLAVES: área urbana, paloma, comportamiento alimentario, ecología de aves, paisaje antropogénico.

INTRODUCTION

The species *Quiscalus mexicanus* (Passeriformes: Icteridae) is an organism strongly associated with anthropogenic places, a highly efficient way of tracking food in urban landscapes. They have sexual dimorphism, males are blue-black, and females are brown with long tails (Christensen 2000, Wehtje 2003, Beedy & Pandolfino 2013). This species is native to Central America and northern South America; however, it is distributed throughout the American continent from Canada to Peru and in some islands of the Caribbean, its affinity for human communities has led it to be a tolerant and adaptable organism (AOU 1998, Christensen 2000, Wehtje 2003, Gurrola-Hidalgo et al. 2009).

In their diet, a variety of foods are known such as fruits, grains, invertebrates such as those found dead on car hoods (Grabrucker & Grabrucker 2010), crabs (*Leptuca beebei*) to small vertebrates. Table 2 provides a reported list of vertebrate preys of *Q. mexicanus*. About predation on rodents, there are only two observational studies. Johnson & Peer (2001) reported for the first time the predation of *M. musculus* and *R. norvegicus* by these birds in North America and Pitts (2020) who describes how a *Q. mexicanus* tried to capture a *Peromyscus maniculatus* in Texas.

Studies related to the predation and feeding mode of birds contribute information on behavior in foraging areas, type of food, and different ecological aspects (Montevecchi 1993, Gill 1995, Barrett et al. 2007, Karnovsky et al. 2012). Different techniques are used to acquire information without disturbing the behavior of birds, such as the use of

telescopes and photographs (Leary 2004, Larson & Craig 2006). Here we report the predation of *Mus musculus* and *Rattus norvegicus* (Rodentia: Muridae), *Columbina talpacoti* (Aves; Columbiformes) and a crustacean of the genus *Leptuca* sp by females of *Q. mexicanus* (Table 1.) to expand the knowledge about its diet and promote research on the changes that *Q. mexicanus* makes in its diet depending on the availability of nutritional resources.

RECORDS.

Observations reported here were not planned, they just occurred by chance in different dates and localities of the province of Panama as reported in Table 1. A Panasonic camera, model Lumix DMC-FZ8 and a Sony camera, model ILCE-7RM2 v4.0. were employed to take pictures and support this report. Observations were made between 7:00 and 13:00 h. Each predation event was observed only one time.

DISCUSSION

Table 1. List of predated animals.

Preys (dam)	Location (coordinates)	Male or female (predator)	Date	Time (hour)
<i>R. norvegicus</i>	09°05'16.7" N; 79°17'10.8" W	Female	13/4/2008	7:06
<i>M. musculus</i>	08°57'48.0"N; 79°32'14.4" W	Female	10/7/2020	12:33
<i>M. musculus</i>	08°59'35"N; 79°32'5" W	Female	30/7/2020	7:43

<i>C. talcopati</i>	8°59'14.2"N	Female	17/8/2020	9:00
	79°31'12.3"W			
Crutacea	9°00'21.4"N	Female	5/9/2020	6:40
(crab)	79°29'08.0"W			

Both preys of Muridae were predated using similar strategies. Males and females of *Q. mexicanus* were observed watching the rodents from a power line but only females flew quickly over them, captured the prey, and pecked them to death. The preys were taken by the neck and strangled breaking their cervical vertebrae. This coincides with the data from Cupul-Magaña et al (2018). The specimen of *C. talpacoti* was captured by a female of *Q. mexicanus* while it was feeding on the ground, it was pecked on its head until died then parts of its body were ripped open and used for food. A female *Q. mexicanus* was photographed while it was carrying in its peak a specimen of *Leptuca* sp. Probably this common crab was captured while was out of its burrow. These reports on food resources of *Q. mexicanus* provide information of the role of *Q. mexicanus* in the community, food web and productivity of their foraging areas, as well as their nutritional requirements for survival and reproduction (Karnovsky et al. 2012).

Despite *Q. mexicanus* is a successful species shown by its wide distribution through the American Continent, studies on its behavior and biology are limited. In its diet are included invertebrates and vertebrates as well as grain and fruits. Table 2 provide information on vertebrate species reported in the literature. Vertebrate's preys are mostly represented by lizards followed by two species of birds and rodents (Sibley, 2001; Gurrola-Hidalgo et al. 2009). Unfortunately, there are not reports available on the invertebrate species of its diet.

The two species of rodents reported here are abundant in the city of Panama, especially in areas where household garbage accumulates, so they can be a food resource that is easily accessible and available all year round. *Q. mexicanus* finds in these garbage accumulations not only

rodents and other vertebrates that supply their protein requirements but also a great diversity of discarded human food.

This report provides information on the food resource of *Q. mexicanus* in Panama and highlight how anthropogenic factors such as garbage accumulation becomes a source of food for them. In this way, we can state that these birds are beneficial as natural controllers for human disease vectors such as rodents. Since *Q. mexicanus* is still expanding its geographical distribution (Johnson and Peer 2001), future studies are necessary to provide a more complete list of vertebrate and invertebrate preys. In addition, it is necessary to determine what environmental or biological factors are involved in their prey selection as well as their impact on the structure of the local rodent population and other vertebrates such as reptiles and birds. The feeding habits of numerous species of neotropical birds have been poorly documented, and this is one of the first records of *Q. mexicanus* predation in Panama. The relevance of this report lies in the information it presents on the feeding behavior and food resources of *Q. mexicanus* in the city of Panama. Perhaps humans must start to consider *Q. mexicanus* as a potential ally in the control of vectors such as rodents and not as a nuisance.

Table 2. List of vertebrate species reported as preys of *Q. mexicanus*.

Class Pisces	No report was found, although there are pictures on the internet.
Class Amphibia	<i>Lithobates neovolcanicus</i> (Cruz-Sáenz et al. 2020)
	Anuran tadpoles (Cupul-Magaña et al 2018)
Class Reptilia	Common house gecko <i>Hemidactylus frenatus</i> (Sánchez-Soto 2015; de Luna et al 2020)
	<i>Anolis</i> sp (Teather and Weatherhead 1988)
	Hatchlings of sea turtle <i>Lepidochelys olivacea</i> (Cupul-Magaña et al 2018)
	Lizards of <i>Aspidozelis sonorae</i> , <i>Sceloporus occidentalis</i> <i>Anolis sagrei</i> (Cupul-Magaña et al 2018)
	<i>Sceloporus melanorhinus</i> , <i>Iguana iguana</i> (Cupul-Magaña et al 2018)
Class Aves	Barn swallow <i>Hirundo rustica</i> . Chicks of the Asian white-winged pigeon <i>Zenaida asiatica</i> (Gurrola-Hidalgo et al. 2009)
Class Mammalia	<i>M. musculus</i> <i>R. norvegicus</i> (Johnson & Peer 2001)




Fig. 1. *Q. mexicanus* feeding on; A. *Leptuca*, B. *M. musculus*, C. *R. novergicus*, D. *M. musculus*.

AKNOWLEDGMENTS

The authors thank the two reviewers, Dr. Daniel Cruz-Saénez and Dr. David Lazcano for their valuable comments that enriched this work.

REFERENCES

AOU: American Ornithologists' Union (1998) Checklist of North American Birds. 7th ed. American Ornithologists' Union Washington, DC.

Barrett, RT, KCJ Camphuysen, T Anker-Nilssen, JW Chardine, RW Furness, S Garthe, O Hüppop, MF Leopold, WA Montevecchi, RR Veit (2007) Diet studies of seabirds: a review and recommendations. ICES Journal of Marine Sciences 64:1675-1691.

Beedy, E. C. & Pandolfino, E. R. (2013). Blackbirds and Relatives Family Icteridae. University of California Press, 335-336.

Christensen, AF (2000) The fifteenth- and twentieth-century colonization of the Basin of Mexico by the Great-tailed Grackle (*Quiscalus mexicanus*). Global Ecology and Biogeography 9:415-420.

Cruz-Saénez, D, D Lazcano, JO Ríos-Martínez, JA García-Salas, A Rodríguez-López & LD Wilson (2020). Notes on the Herpetofauna of Western Mexico 24: Predation on *Lithobates neovolcanicus* (Hillis & Frost, 1985) by *Quiscalus mexicanus* (Gmelin, 1788) in the Gardens of Guadalajara Zoo, Jalisco, Mexico. Bulletin of the Chicago Herpetological Society 55: 153-157.

Cupul-Magaña, FG, F Mc Cann & AH Escobedo-Galván (2018). Observaciones generales de la dieta del zanate mexicano *Quiscalus mexicanus* en Puerto Vallarta, México. Huitzil, 19: 96-99.

de Luna, M., Rodríguez, M., & Barrios, R. G. (2020). Predation on the common house gecko *Hemidactylus frenatus* Schlegel, 1836 by the brown widow spider *Latrodectus geometricus* Koch, 1941 in Colima, Mexico. Herpetology Notes, 13, 555-556.

Gill, FB (1995) Ornithology. W.H. Freeman and Company, Nueva York, E.U.A.

Grabruker, S, AM Grabruker (2010). Rare feeding behavior of Great-Tailed Grackles (*Quiscalus mexicanus*) in the extreme habitat of Death Valley. *The Open Ornithology Journal* 3:101-104.

Gurrola-Hidalgo, MA, C Sánchez-Hernández, ML Romero-Almaraz (2009) Novel food sources for *Quiscalus mexicanus* and *Cyanocorax sanblasianus* in Chamela, Jalisco coast, Mexico. *Acta Zoológica Mexicana* 25:427-430.

Johnson, K & BD Peer (2001). Great-tailed Grackle (*Quiscalus mexicanus*). Pp. 1-28. En A. Poole, F. Gill (eds.), *The Birds of North America* 576. The Birds of North America, Inc., Philadelphia, E.U.A.

Karnovsky, NJ, KA Hobson, SJ Iverson (2012) From lavage to lipids: estimating diets of seabirds. *Marine Ecology Progress Series* 451: 263-284.

Larson, K, D Craig (2006). Digiscoping vouchers for diet studies in bill-load holding birds. *Waterbirds: The International Journal of Waterbird Biology* 29: 198-202.

Leary, PR (2004) Digiscope applications for shorebird studies. *Wader Study Group Bulletin* 104: 34-38.

Montevocchi, WA (1993). Birds as indicators of change in marine prey stocks. Pp. 217-266. En R.W. Furness, J.J.D. Greenwood (eds.), *Birds as monitors of environmental change*. Chapman & Hall, Londres, Reino Unido.

Pitts, RM (2020). Note of the attempted predation of an adult deer mouse (*Peromyscus maniculatus*) BY A GREAT-TAILED GRACKLE (*Quiscalus mexicanus*). *Transactions of the Missouri Academy of Science* 48: 4-4.

Sánchez-Soto, S (2015). Depredación de *Hemidactylus frenatus* (Reptilia) por *Quiscalus mexicanus* (Aves). *Zeledonia* 19: 125-127.

Sibley, C. 2001. The Cornell lab of ornithology's guide to birds of North America version 3.0 (Electronic media- CD). Thayer Birding Software.
Teather, K. L. and P. J. Weatherhead. (1988). Sex-specific energy requirements of Great-tailed Grackle (*Quiscalus mexicanus*) nestlings. Journal of Animal Ecology 57:659-668.

Wethje, W (2003) The range expansion of the great-tailed grackle (*Quiscalus mexicanus* Gmelin) in North America since 1880. Journal of Biogeography 30: 1593-1607.

Recibido 20 diciembre 2020, y aceptado el 13 mayo 2021
Editor Responsable: Dr. Alonso Santos Murgas